

**CHECKLIST OF THE AQUATIC MACROPHYTES IN
CERTAIN WETLANDS OF YAVATMAL DISTRICT,
MAHARASTRA STATE, INDIA**

**MUKUND DHORE^{1*}, MANIK DHORE², PARESH LACHURE¹
AND DINESH DABHADKAR³**

¹Department of Botany, B. B. Arts, N. B. Commerce & B. P. Science College,
Digras, Maharashtra, India - 445203

²Shivaji College, Akot. Dist.- Akola, Maharashtra, India

³Department of Zoology, Govt. Vidarbha Institute of Science and Humanities,
Amravati, Maharashtra, India- 444604

ABSTRACT

The present investigation concerns with the aquatic macrophyte diversity and its role in understanding the wetland ecosystem dynamics and species composition of aquatic macrophytes, seasonal distribution in five wetlands in Yavatmal district, viz. Arunavati dam, Nandgavan dam, Fetri dam in Digras Tehsil and Jamwadi Lake and Manpur Lake, in Yavatmal tehsil were studied during 2009-2011. Forty two different species of aquatic macrophytes were recorded from the studied wetlands which include three free floating macrophytes, seven submerged, two rooted floating and twenty eight emergent species of macrophytes. The free floating species *Eichornia crassipes* occurs throughout the year. While the submerged species *Vallisneria spiralis*, *Ceratophyllum demersum*, *Hydrilla verticillata* occur throughout the year. In the rooted floating category *Ipomoea aquatica* occurs throughout the year.

**Checklist of the Aquatic Macrophytes in Certain Wetlands of
Yavatmal District, Maharashtra State, India**

Species composition and seasonal distribution of the macrophytes has also been noted in the present paper.

KEY WORDS: Aquatic macrophytes, Yavatmal District, Digras tehsil, Wetlands.

INTRODUCTION

Macrophytes serve as a link between the sediment, water, and (sometimes) atmosphere in wetlands, lakes, and rivers. The most notable function that plants serve is as primary producers. However, macrophytes are also involved in ecosystem processes such as biomineralization, transpiration, sedimentation, elemental cycling, materials transformation, and release of biogenic trace gases into the atmosphere (Carpenter and Lodge, 1986). Recent studies also suggest that macrophytes play a central role in shallow lakes which can have two possible stable equilibria: a clear-water state that is dominated by aquatic macrophytes and a turbid-water state that is dominated by phytoplankton (Scheffer *et al.*, 1993; Moss *et al.*, 1994, Jeppesen *et al.*, 1998). Macrophytes maintain the clear-water state by a variety of mechanisms whose relative importance is probably variable (Ozimek *et al.*, 1990, Vermaat *et al.*, 2000, Madsen *et al.*, 2001). Aquatic weeds referred to as Macrophytes constitute an important component of aquatic ecosystem. Their diversity and biomass influence primary productivity and complexities of trophic states (Kumar and Singh, 1987).

Since the wetland is considered as a transitional area between land and water, Smith (1980) aptly described it as a half-way world between terrestrial and aquatic ecosystems. It is largely dominated by water and has special type of flora and fauna, which usually undergo time scheduled characteristic changes from hydric to mesic types. Wetlands are well known for high diversity in class, composition and four broad categories of functions *viz.* physical, hydrological, chemical, biological and socioeconomic (Williams, 1990). Wetland supports

**Mukund Dhore, Manik Dhore, Paresh Lachure
And Dinesh Dabhadkar**

plant species intermediate between true aquatic and terrestrial habitats (Banerjee and Venu, 1994).

Thus researches on wetland macrophytes have started gaining importance not only because systematic stock taking of biodiversity is presently given top most priority but also because these plants have implications with functional values of wetlands. In addition to stock taking and assessment of ecological functions of different forms, wetlands biodiversity pattern is studied emphatically. Vegetation pattern are likely to control major aspects of wetlands biogeochemistry and tropic dynamics and wetlands should be viewed as complex mosaic of habitats with distinct structural and functional characteristics (Rose and Crompton, 1996). There is very scanty literature available about the aquatic macrophytes. The present investigation was, therefore, undertaken to study the species composition and seasonal distribution of aquatic macrophytes in certain wetlands of Yavatmal district, Maharashtra.

METHODS AND MATERIAL

The study area i.e. Yavatmal district is situated in the eastern part of Maharashtra between north latitudes $19^{\circ} 28'$ and $20^{\circ} 48'$ and East longitudes $77^{\circ} 18'$ and $79^{\circ} 98'$. This region is blessed with a good number of fresh water lakes and dams harbouring a great variety of aquatic macrophytes. Information on phytosociological data for aquatic macrophytes in any water body is of immense importance to understand the wetland ecosystem. Much work has been done on the phytosociology of different macrophytic species in different freshwater bodies of India and abroad (Billore and vyas, 1981, Biswas and Calder, 1984). In the present study monthly survey was done by quadrat method for collecting submerged aquatic macrophytes from January 2009 to July 2011. Qualitative and quantitative analysis of aquatic macrophytes was done by following the methodology of Biswas and Calder (1984) and Mishra (1974). In each water

**Checklist of the Aquatic Macrophytes in Certain Wetlands of
Yavatmal District, Maharashtra State, India**

body, the aquatic macrophytes were analyzed and specimens were identified up to genus/species level.

RESULTS

Aquatic macrophytes exhibited a heterogenous assemblage of forty one species in the studied wetlands. These species were distributed in thirty eight genera representing a variety of taxonomic group angiosperms, charophyta and bryophyta. Out of these *Eichornia crassipes*, *Vallisneria spiralis*, *Hydrilla verticillata*, *Ipomoea aquatica*, *Cynodon dactylon* occur throughout the year. During monsoon, some floating species of aquatic macrophytes were found like *Eichornia crassipes*, *Vallisneria spiralis*, *Hydrilla verticillata*, *Ipomoea aquatic* which flourished and dominated the aqua-regime at a greater depth of water. Conversely, the monsoon varieties of aquatic macrophytes were succeeded by the winter emergent species viz. *Lemna perpusilla*, *Limnophilla sessiflora*, *Ottelia alismoides*, *Najas minor*. The species that occur during the onset of summer were observed to be *Heliotropium supinum*, *Crozophora rotleri*, *Gnaphalium pulvinatum*, *Glinus lotoides*. Significant phytosocial associations have been recorded among the different aquatic macrophytes like *Hydrilla verticillata*, *Vallisneria spiralis*, *chara sp.*, *Nitella sp.* Similarly *Vallisneria spiralis*, *Najas minor*, *Potamogeton pectinatus*, *Ceratophyllum demersum* and *Ottelia allismoides*, *Limnophilla sessiflora*, *Valisneria spiralis* were also found to be in associated with each other.

DISCUSSIONS

The check list generated in the study is intended to support other research in wetlands and in particular, to assure the continuity of ongoing long term ecological programs. Cowardin *et al.*, (1979) gave an idea on the checklist of wetlands and deep water habitats of macrophytes of the United States. Keddy (2000) classified macrophytes on the basis of wetlands habitat of macrophytes.

**Mukund Dhore, Manik Dhore, Paresh Lachure
And Dinesh Dabhadkar**

The aquatic macrophytes are classified as submerged, floating and emergent growing in and covered by at least 25cm of water.

Kayode and Ogunleye (2008) were also studied the checklist and status of plant species used as spices in Kaduna State of Nigeria. *Eichhornia crassipes*, which is listed in 1995 as one of the invasive, problematic aquatic plants. Cronk and Fuller, 1995 was found also recorded the same type of checklist. The prominent macrophytes recorded during this study were *V. nigritana*, *S. dalzielii* and *T. australis*. *Sesbania dalzielii*, which the riparian communities livestock farmers depend on as a forage plant (Obot, 1984; Adesina *et al.*, 2007) and whose growth is naturally tied to rainy seasons, does not pose threat to the lake. However, the other newly recorded macrophytic species pose major threat to the lake existence. Chambers *et al.*, 2010 were studied the world checklist of macrophyte species.

Thomaz *et al.* (1999) found 62 taxa belonging to 25 families and 42 genera. Among this total, 47 were identified at the species level and 33 species were common between both surveys. Similar to what we found in present study. Differences in the number of species and the identity of genera between these surveys indicate that macrophyte assemblages are still changing and are dynamic in this reservoir.

Checklist of the Aquatic Macrophytes in Certain Wetlands of
Yavatmal District, Maharashtra State, India

**Table 1 : Checklist of some aquatic macrophytes in certain wetlands of
Yavatmal district**

Sl. No.	Name of Species	Family	Life Forms
1	<i>Aponogeton natans</i> L.f	Aponogetonaceae	Submerged
2	<i>Alternanthera sessilis</i> (L.)R. Br. ex DC	Amaranthaceae	Emergent
3	<i>Ammania baccifera</i> L.	Lythraceae	Emergent
4	<i>Aeschonemene indica</i> L.	Fabaceae	Emergent
5	<i>Bacopa monnieri</i> (L.) Wettestin	Scrophulariaceae	Emergent
6	<i>Chara</i> sp.	Characeae	Submerged
7	<i>Ceratophyllum demersum</i> L.	Ceratophyllaceae	Submerged
8	<i>Coix aquatica</i> Roxb.	Poaceae	Emergent
9	<i>Commelina benghalensis</i> L.	Commelinaceae	Emergent
10	<i>Commelina hasskarlii</i> C. Comm. Cyrt.	Commelinaceae	Emergent
11	<i>Cyperus rotundus</i> L.	Cyperaceae	Emergent
12	<i>Cynodon dactylon</i> (L.) Pers.	Poaceae	Emergent
13	<i>Cyathocline purpurea</i> (Buch-Ham. ex D. Don) Oltze	Asteraceae	Emergent
14	<i>Crozophora rottlerii</i> (Geisel.) A. Juss. ex. Spr.	Euphorbiaceae	Emergent
15	<i>Dopartium junecum</i> (Roxb.) Buch-Hum ex Benth.	Scrophulariaceae	Emergent
16	<i>Elaeocharis geniculata</i> (L.) R&S.	Cyperaceae	Emergent
17	<i>Elaeocharis capitata</i> R. Br.	Cyperaceae	Emergent
18	<i>Eriocaulan cinereum</i> R.Br.	Eriocaulaceae	Emergent

Mukund Dhore, Manik Dhore, Paresh Lachure
And Dinesh Dabhadkar

19	<i>Echinochloa calanum</i> (L.) Link	Poaceae	Emergent
20	<i>Eichornia crassipes</i> (Mart.) Solms.	Pontederiaceae	Free floating
21	<i>Gomphrena celosioides</i> Mart.	Amaranthaceae	Emergent
22	<i>Glinus lotoides</i> L.	Molluginaceae	Emergent
23	<i>Gnaphalium pulvinatum</i> Del.	Asteraceae	Emergent
24	<i>Hydrilla verticillata</i> (L. f.) Royle	Hydrocharitaceae	Submerged
25	<i>Heliotropium supinum</i> L.	Boraginaceae	Emergent
26	<i>Ipomoea aquatica</i> Forsk	Convolvulaceae	Rooted floating
27	<i>Lemna perpusilla</i> Torrey	Lemnaceae	Free floating
28	<i>Limnophylla sessiflora</i> L.	Plantaginaceae	Emergent
29	<i>Najas minor</i> L.	Hydrocharitaceae	Submerged
30	<i>Ottellia alismoides</i> (L.) Pers.	Hydrocharitaceae	Rooted floating
31	<i>Pistia stratioides</i> L.	Araceae	Free floating
32	<i>Potamogeton pectinatus</i> L.	Potamogetonaceae	Submerged
33	<i>Polygonum glabrum</i> Willd.	Polygonaceae	Emergent
34	<i>Phyla nodiflora</i> (L.) Greene	Verbinaceae	Emergent
35	<i>Riccia sp.</i>	Ricciaceae	Emergent
36	<i>Rotala serpillifolia</i> (Roth.) Breneck	Lythraceae	Emergent
37	<i>Sesbania bispinosa</i> (Jacq.) w. t. wight	Fabaceae	Emergent
38	<i>Sopubia delphinifolia</i> (L.) G. Don	Scrophulariaceae	Emergent
39	<i>Sphaeranthus indicus</i> L.	Asteraceae	Emergent
40	<i>Typha angustata</i> Bory and Chaub.	Typhaceae	Emergent
41	<i>Vallisneria spiralis</i> L.	Hydrocharitaceae	Submerged
42	<i>Verbascum chinense</i> (L.) Sant	Scrophulariaceae	Emergent

**Checklist of the Aquatic Macrophytes in Certain Wetlands of
Yavatmal District, Maharashtra State, India**

REFERENCES

1. Adesina, G. O., Akinyemiju, O.A. and Olaleye, V.F. (2007) Assessment of aquatic vegetation of Jebba Lake, Nigeria. *African Journal of Ecology* 45:365-373.
2. Banerjee, L. K. and Venu, P. (1994) Wetlands plant resources for conservation. E&VIS (Newsletter of Botanical Survey of India, Calcutta) 1: 2-3.
3. Billore, D. K. and Vyas, I. N. (1981) Distribution and production of macrophytes in pichhola lake, Udaipur. *Dnt J Ecol Env-sci* 7:45-54.
4. Biswas, K. and Calder, L. C. (1984) Handbook of common water and marsh plants of India and Burma, xvii + 216, Bishensingh Mahendra palsingh [Dehradun].
5. Carpenter, S. R., Lodge, D. M. (1986) Effects of submersed macrophytes on ecosystem processes. *Aquatic Bot* 26: 341-370.
6. Chambers, P.A., P. Lacoul, K.J., Murphy, S.M., Thomaz and Z. Duggan.(2010) World checklist of macrophyte species. Published on the internet; <http://fada.biodiversity.be/group/show/60> accessed 10 August 2010.
7. Cowardin et al., (1979)
8. Cronk, Q.C.B. and Fuller, J. L. (1995) *Plant invaders: the threat to natural ecosystems*. Chapman and Hall, London. 241pp.
9. Jeppesen, E., Søndergaard, M., Søndergaard, M., Christoffersen, K. (Eds.). (1998) The Structuring Role of Submerged Macrophytes in Lakes. *Ecological Series*, vol. 131. Springer-Verlag, 423 pp.
10. Kayode, J. and Ogunleye, O.T. (2008) Checklist and Status of Plant Species Used as Spices in Kaduna State of Nigeria. *African Journal of General Agriculture* 4, 13-18.

**Mukund Dhore, Manik Dhore, Paresh Lachure
And Dinesh Dabhadkar**

11. Keddy (2000)
12. Kumar, M. and Singh, J. (1987) Environmental impacts of Aquatic Weeds and their classification. Proceedings of the workshop on management of Aquatic Weeds, Amritsar, Punjab, India.
13. Madsen, J.D., Chambers, P.A., James, W.F., Koch, E.W., Westlake, D.F. (2001) The interaction between water movement, sediment dynamics and submersed macrophytes. *Hydrobiologia* 444: 71-84.
14. Mishra, K.C. (1974) Manual of plant ecology, *oxford and IBH publishing co.* New Delhi 491.
15. Moss, B., McGowan, S., Carvalho, L. (1994) Determinations of phytoplankton crops by top-down and bottom-up mechanisms in a group of English lakes, the West Midland meres. *Limnol Oceanogr* 39: 1020-1029.
16. Obot, E.A. (1984) Studies on the productivity of *Echinochloa stagnina*(Retz) P. Beauv in the Kainji Lake Basin of Northern Nigeria *Ph.D. Thesis, University of Ife, Nigeria.*
17. Ozimek, T., van Donk, E., Gulati, R.D., (1990) Can macrophytes be useful in biomanipulations of lakes? The Lake Zwemlust example. *Hydrobiologia* 201:399-409.
18. Rose, C. and Crumpton, W.G. (1996) Effects of emergent macrophytes on dissolved oxygen dynamics in a prairie pothole wetland. *Wetland. Ecology* 19: 105-115.
19. Scheffer, M., Hosper, S.H., Meijer, M.L., Moss, B., Jeppesen, E. (1993) Alternative equilibria in shallow lakes. *Trends Eco. Evol* 8: 275-279.
20. Smith, R. I. (1980) *Ecology and Field Biology* (3rd .Ed) Harper and Row. New York.
21. Thomaz, S.M., L.M. Bini, M.C., Souza, K.K. Kita and Camargo. (1999) Aquatic Macrophytes of Itaipu Reservoir, Brazil: Survey of Species and Ecological Considerations. *Braz Arch Biol. Techn* 42: 15-22.

**Checklist of the Aquatic Macrophytes in Certain Wetlands of
Yavatmal District, Maharashtra State, India**

22. Vermaat, J.E., Santamaria, L., Roos, P.J. (2000) Water flow across and sediment trapping in submerged macrophyte beds of contrasting growth form. *Arch Hydrobiol* 148:549-562.
23. Williams, M. (1990) Understanding Wetlands in Michael Williams (ed), *Wetlands: A threatened Landscape. Basil Blackwell Ltd., Oxford, 1990.* pp.1-3.
24. Cowardin, L. M., Carter.